



# First ethnomycological record of *Fistulinella wolfeana* as an edible species and some of its nutritional values

## Primer registro etnomicológico de *Fistulinella wolfeana* como una especie comestible y algunos valores nutricionales

Daniel Robles-García <sup>1</sup>, Elhadi Yahia <sup>2</sup>, Jesús García-Jiménez <sup>3</sup>, Edgardo Ulises Esquivel-Naranjo <sup>4</sup>, Fidel Landeros <sup>1</sup>

<sup>1</sup>Laboratorio de Ecología y Sistemática de Microorganismos, Facultad de Ciencias Naturales, Universidad Autónoma de Querétaro, México. <sup>2</sup>Laboratorio de Fitoquímicos y Nutrición Humana, Facultad de Ciencias Naturales, Universidad Autónoma de Querétaro, México. <sup>3</sup>Instituto Tecnológico de Ciudad Victoria, México. <sup>4</sup>Laboratorio de Microbiología Molecular (LAMIMO), Facultad de Ciencias Naturales, Universidad Autónoma de Querétaro, México

### RESUMEN

Registramos por primera vez el uso de *Fistulinella wolfeana* como especie comestible en las comunidades otomías de Tesquedó, Xajay y Tenasdá, en Amealco de Bonfil, Querétaro, México, la cual forma parte de los recursos naturales utilizados durante la estación de lluvias. Además, esta especie está ampliamente distribuida en el Eje Volcánico Transversal, y tiene el potencial de ser utilizada como recurso no maderable. Se describe el valor nutricional de las muestras en fresco, así como en seco. Además, tiene una alta actividad antioxidante, alcanzando casi 90% de inhibición en diluciones 1:10 de extracto metanólico, comparable con valores para otros miembros de la familia Boletaceae.

**PALABRAS CLAVE:** hongos comestibles, etnomicología, valor nutritivo, actividad antioxidante.

### ABSTRACT

We registered for first time the use of *Fistulinella wolfeana* as an edible species in the Otomi communities of Tesquedó, Xajay and Tenasdá, in Amealco de Bonfil, Querétaro, México, which is part of the natural resources used during the rainy season. In addition, this species is widely distributed in the Volcanic Transmexican Belt, and it has the potential to be used as non-timber resource. The nutritional value for fresh, as well as dried samples, is described. Additionally, it has high antioxidant activity, reaching almost 90% of inhibition in 1:10 dilutions of methanolic extract, comparable with values for other members of the Boletaceae family.

**KEYWORDS:** edible mushroom, ethnomycology, nutritional value, antioxidant activity.

### INTRODUCTION

Mexico is a high diversity center of the Boletaceae family, with 212 known taxa (García and Garza-Ocañas, 2001) of 877 species described in the world (Kirk *et al.*, 2008). *Fistulinella wolfeana* Singer & J. García was described as a Mexican new

species (Singer *et al.*, 1991), however, its edibility was not determined. González and Valenzuela (1993) developed an inventory of Boletaceae and Gomphidiaceae from the State de México, and reported their seasonality and distribution, but they did not

Recibido / Received: 19/02/2016

Aceptado / Accepted: 30/11/2016

Autor para correspondencia / Corresponding author:

Fidel Landeros

landeros@uaq.mx

mention them as an edible species. This is a very common species in Mexican forests (García *et al.*, 1998; García, 1999; García and Garza-Ocañas, 2001). Villanueva *et al.* (2008) mentioned *F. wolfeana* as a mycorrhizal species with *Quercus* spp., and García (1999) described its distribution along the Mexican central states, such as Jalisco, Guanajuato, Querétaro, Morelos, the State of México and Hidalgo, but it could be found in other states.

Gomphidiaceae from the State de México, and reported their seasonality and distribution, but they did not mention them as an edible species. This is a very common species in Mexican forests (García *et al.*, 1998; García, 1999; García and Garza-Ocañas, 2001). Villanueva *et al.* (2008) mentioned *F. wolfeana* as a mycorrhizal species with *Quercus* spp., and García (1999) described its distribution along the Mexican central states, such as Jalisco, Guanajuato, Querétaro, Morelos, the State of México and Hidalgo, but it could be found in other states.

Estrada-Torres and Aroche (1987), Burrola-Aguilar *et al.* (2012) and Lara-Vázquez *et al.* (2013) studied the ethnomycological knowledge of some Otomí communities in State of México and reported 28 edible mushrooms species. At the national level, Queretaro State is the third largest Otomí population (INEGI, 2009), however, there has been no reported edible mus-

hrooms species related to this ethnic group. There is only evidence that two species are consumed in Querétaro, particularly near the “Cerro el Zamorano” in the municipality of Colón (Landeros *et al.*, 2006), *Clavulina amethystina* named “menudo de vaca” and *Lycoperdon perlatum* as “huevo” (egg).

The aim of the present work is to report the edibility of *F. wolfeana* for the first time, the ethnomycological importance as a non-timber forest product, and its use as a food resource, with a general description of its nutritional value and antioxidant capacity.

## MATERIALS AND METHODS

### Collected material

The specimens were bought in Tesquedó community, and were used to determine the species taxonomic keys of Singer *et al.* (1991). About 700 g of fresh mushroom were collected from Xajay community (a locality near to Tesquedó), covered with aluminum foil, placed inside a plastic bag and transported to the laboratory in a cooler with refrigerant.

### Color determinations

Color determination was made on pileus, context and pores, of three fresh mushrooms of *F. wolfeana*. Measurements were




Figure 1. Location of the study area.



taken five times per structure (pileus, context and pores) with a Minolta spectrophotometer (CM-2002). The parameters  $L^*$  (Luminosity),  $a^*$  (red/Green coordinates),  $b^*$  (yellow/blue coordinates),  $c^*$  (chrome and clarity) and  $h^\circ$  (tone angle) were averaged and compared using standard deviation. For pore surface and context, precautions were taken to avoid any damage that could induce color changes, caused by touching these parts. The colors are important since wild mushrooms are an important source of bioactive compounds, such as pigments (Ferreira *et al.*, 2009).

### Proximal analysis

One sample of 300 g of fresh mushrooms and one sample of 300 g of dried mushrooms were processed to determine their nutritional values. The proximal analysis was conducted in the Unit of Chemical Services at the Faculty of Chemistry of the Universidad Autónoma de Querétaro, using the methods established by the Mexican norms: NMX-F-607-NORMEX-2002) for ash determination, NMX-F-613-NORMEX-2003 for raw fiber determination, NMX-F-608-NORMEX-2002 for protein determination, NMX-F-089-S-1978 for ethereal extracts determinations (Soxhlet method), and NMX-AA-051-SCFI-2001 for the use of atomic absorption spectrophotometer (Perkin Elmer, 2002).

### Antioxidant activity

Mushroom extracts were prepared according to Elmastas *et al.* (2007) with some modifications. For methanolic extraction, 1 g of powdered lyophilized mushrooms was resuspended in 40 mL of methanol and then mixed with a homogenizer (IKA T25 Basic) into Falcon tubes, and were covered with aluminum foil and placed on an orbital shaker (Orbital shaker Midi OM10E) for 24 h. The extract was filtered with a Whatman no. 2 paper and the remaining extract was collected. Methyl alcohol was removed in a rotatory evaporator at 40°C and 70 rpm at 175 mbar to obtain the dried extract (Büchi, Heating Bath B490, Rotovapor R-205 and Vacuum V-800). The final amount of the extract was resuspended in 2 mL of HPLC water and stored at 4°C for one day, to avoid oxidative damage.

The radical scavenging activity was determined with the 2,2-diphenyl-1-picrylhydrazyl (DPPH) assay, as reported by Brand-Williams *et al.* (1995), and was modified to read in a

96-well microplate. Aliquots of 280  $\mu$ L of DPPH (100  $\mu$ M) in methanol were deposited with 20  $\mu$ L of mushroom extract at 1:10 and 1:100, and completed to 300  $\mu$ L each. Samples were incubated in the dark for 45 min, and then measured at 490 nm (Microplate reader). Samples were processed in triplicate. Trolox reagent was used as standard, and was prepared at 0, 0.2, 0.4, 0.5, 0.6, 0.7, 0.8 and 1  $\mu$ M to determine Trolox equivalents.

## RESULTS AND DISCUSSION

### Ethnomycological importance

*Fistulinella wolfeana* was found in local markets for the first time in the Tesquedó community, and then was mentioned by several people who live in the three communities (Tesquedó, Xajay and Tenasdá). It is known as “Ushki jieth’é” in otomí dialect, meaning “salted mushroom”, because people attribute it a salty taste, even though some call it “sweet mushroom”, due



Figure 2. People from Tesquedó community who collect *Fistulinella wolfeana* for sale.

to a mild flavor when the mushroom is mature. It is collected either for self-consumption, economic purposes (Figure 2), or even to exchange for main supplies, such as vegetable oil, bread, rice or corn. This is important to emphasize, because there are no previous reports of its edibility. Therefore, *F. wolfeana* could be used both as a food source or an economical income source. An important feature is that people dry and keep the mushrooms (Figure 3) for almost a year, to consume them during dry season, when there are not mushrooms and food is scarce. The preservation is possible because this species is rarely invaded by larvae, even when it is left on the soil exposed to sunlight. In fact, this is the first report of conservation of a bolete, because *Boletus* spp., and other genera of the Boletaceae family, are fre-

quently infected by larvae, therefore, people cannot conserve them. Estrada-Torres and Aroche (1987) indicated that people from Acambay, Estado de México, dry their mushrooms hanging them with a thread, and exposing them to the sunlight; but there was no mention of any bolete among them. In our study, we have observed that people place the mushrooms over a rock or a tree during the day, and at night they keep them inside to avoid rehydration, and repeat the same process for few days until the mushrooms are completely dry. During this study, *F. wolfeana* was consumed for two consecutive seasons by the authors and at least seven more people, who indicated that the taste was exquisite.

## SPECIES DESCRIPTION

*Fistulinella wolfeana* Singer & J. García, in Singer, García & Gómez, Beih. Nova Hedwigia 102: 82 (1991)

Figure 4

Pileus 30-150 mm, plane-convex, plane-concave or concave, brown pinkish to violet, glabrous; pores 1-1.5 mm diameter, rounded, pinkish, orange-reddish when it is injured, sometimes emarginated; individual tubes 7-20 mm long, white becoming pinkish in older specimens; stipe cylindrical to bulbous at the base; context 15-50 mm, mostly white, unchanging or slightly pinkish in the pileus, yellowish to the base of the stipe, and taste of pileus is mild.

Spores 10-14 × 4-5.6 μm, yellowish to ochraceous yellow, subfusoid, some with suprahilar depression; basidia 23-28 × 9-11 μm, 4-spored; pleurocystidia 36-55 × 6-12 μm, mostly ventric to fusiform, hyaline to yellowish in KOH and reddish in Melzer's reagent; cheilocystidia 38-60 × 5-12 μm, ventric to fusiform mostly, ochraceous; hymenophoral trama bilateral, with the center gelatinized, slightly ochraceous; epicutis of the pileus with hyphae 2-7 μm diam., composed of a gelatinized ixotrichodermium, with parallel elements cystidia like, with rounded apex mostly yellowish with ochraceous lipid content; cells of the stipe 21-30 × 4-6(-7) μm, versiform, with cystidia like elements and dermatopseudocystidia, 4-spored caulobasidia and terminal elements with rounded apex.



Figure 3. Dry specimens of sun-dried *Fistulinella wolfeana*.




Figure 4. Basidiome (a). Spores (b), Basidia (c), Cheilocystidia (d), Pleurocystidia (e). Scale bar (a) = 5 cm, (b, c, d, e) = 10µm.

**Comments.** *Fistulinella wolfeana* could be confused with some species of *Tylopilus*, because both genera have rose tubes and pinkish pores, which change to orange or ochraceous reddish when touched. *Tylopilus indecisus* has a pale brown slightly reticulated stipe and thin spores (10.5-15.5 × 3.5-4.5). *Tylopilus plumbeoviolaceus* has a very bitter taste and reticulated stipe surface. Finally, *T. subcellulosus* has narrower and shorter spores (9.5-12 × 3.5-4) and bitter taste, compared with *F. wolfeana*. On the other hand, *F. wolfeana* has viscid surface pileus, while all *Tylopilus* species have subtomentosus, glabrous or smooth dry surface.

**Material studied.** *Robles 10*, 11 Aug 2013, (QMEX); *Robles 162*, 15 Sept 2013 (QMEX), Amealco, Querétaro, México.

**Habitat.** In hardwood forest, mainly by *Quercus* spp. with some mixed reforested areas of *Pinus* spp.

### Color determination

There is a lack of reported quantitative color determination for wild mushrooms, and most determinations have been done on cultivated species with industrial purposes (Mohapatra *et al.*, 2010; Qi *et al.*, 2014). We made a complementation of the original color description, with the values obtained in the test. Pileus and pores show similar values, while context presented differences due to its whitish color. According to color scores, lumino-

sity ( $L^*$ ) had the following order: context > pores > pileus; coordinates red/green ( $a^*$ ): pores > pileus > context; and coordinates yellow/blue ( $b^*$ ): pores > pileus > context. On the other hand,  $c^*$  and  $h^\circ$  parameters had the following order, chrome ( $c^*$ ): pores > pileus > context; and tone angle ( $h^\circ$ ): context > pores > pileus. The parameter in the pileus had the major deviation in color, including pinkish, pinkish-violet or pinkish-brownish, probably due to sun exposure. Pores change their color, from white to pinkish-lilac, when touched. It could explain the minimal variation and similarity between the values of both parts. The high value in context explains the white color and the low variation in color change, as indicated by Singer *et al.* (1991) in the original description. The above values, suggest a variation depending on where the mushroom is growing, and if it is directly influenced by the sunlight, because specimens found in the inner part of the forest had lower  $L^*$  values, indicating a higher color tone, while those exposed to sunlight had paler tones, showing higher  $L^*$  values. Pileus and pores had similar values; context does not change since it is inside, and only pileus shows a considerable variation (Table 1). Thus, this difference may suggest a variation in the amount of pigments that decrease in exposed specimens. It would be convenient to determine pigments, to find out which are the most important that make up the colors of this species, and how they are modified by environmental conditions.

Table 1. Color values of pileus, pores surface and context of *Fistulinella wolfeana*

Parameters	Pileus	Pores	Context
L*	46.37 ± 5.4	49.26 ± 0.9	74.62 ± 0.8
a*	8.62 ± 0.9	8.90 ± 0.5	0.88 ± 0.4
b*	25.09 ± 1.1	25.53 ± 1.7	16.67 ± 0.8
c*	26.74 ± 1.3	27.27 ± 1.9	16.71 ± 0.8
h°	70.24 ± 0.8	70.64 ± 0.9	87.18 ± 1.1

L\* (Luminosity), a\* (red/green coordinates), b\* (yellow/blue coordinates), c\* (chrome and clarity) and h° (tone angle). Values averaged ± standard deviation.

### Proximal analysis

Results of proximal analysis are shown in Table 2, which are compared with data from other foods, such as mushrooms and some meats frequently consumed. Protein levels, in dry weight,

were similar to those in *Agaricus bisporus* and *Pleurotus ostreatus*, but with less than half of fat and carbohydrates. Therefore, *F. wolfeana* is a food with high protein content.

In addition, its sodium content, was very low compared to another mushrooms species. This is important because of potential health problems, such as hypertension that can be caused by excess of sodium. Finally, *F. wolfeana* had more fiber content, in both fresh and dry weight, than any of the commercial mushrooms compared, which means that it can help lower cholesterol levels and promote digestion.

Regarding to comparison with some meats of frequent consumption, protein values (in dry weight) were similar to those in chicken, beef, tuna fish and blacktip shark. *Fistulinella wolfeana* had fewer carbohydrates and fats than meat; such as the case with commercial mushrooms. Two important aspects to consider in *F. wolfeana*, the presence of fiber and the low amount of sodium, indicate that this species is a good nutritional alternative, even compared to meat.

Table 2. Comparison of fresh and dry proximal analysis of *Fistulinella wolfeana* with other foods

Species/Food		Carbohydrates	Ash	Fiber	Fats	Moisture	Protein	Sodium (mg/100g)
<i>Fistulinella wolfeana</i>	Fresh	18.01	1.49	2.93	0.46	73.69	3.42	-
	Dry	30.21	4.48	23	0.71	11.77	29.83	4.5
<i>Boletus gruop</i> dried	Fresh	N	N	N	N	N	N	N
	Dry	19.46 <sup>5</sup>	1.71 <sup>5</sup>	8.74 <sup>5</sup>	1.08 <sup>5</sup>	N	0.12 <sup>5</sup>	N
<i>Agaricus bisporus</i>	Fresh	0.4 <sup>4</sup>	0.82 <sup>5</sup>	2.3 <sup>4</sup>	0.3 <sup>4</sup>	78.3-90.5 <sup>2</sup>	1.8 <sup>4</sup>	21 <sup>3</sup>
	Dry	51.3-62.5 <sup>2</sup>	7.7-12 <sup>2</sup>	8-10.4 <sup>2</sup>	1.7-8 <sup>2</sup>	N	23.9-34.8 <sup>2</sup>	N
<i>Pleurotus ostreatus</i>	Fresh	Trace <sup>4</sup>	0.89 <sup>5</sup>	N	0.2 <sup>4</sup>	73.7-90.8 <sup>2</sup>	1.6 <sup>4</sup>	6 <sup>3</sup>
	Dry	57.6-81.1 <sup>2</sup>	6.1-9.8 <sup>2</sup>	7.5-8.7 <sup>2</sup>	1.6-2.2 <sup>2</sup>	N	10.5-30.4 <sup>2</sup>	N
Chicken	Fresh	25.1 <sup>1</sup>	1.28 <sup>1</sup>	N	7.75 <sup>1</sup>	74.9 <sup>1</sup>	19.7 <sup>1</sup>	51.55 <sup>1</sup>
	Dry	N	N	N	N	N	N	N
Cow	Fresh	27 <sup>1</sup>	0.97 <sup>1</sup>	N	2.57 <sup>1</sup>	73 <sup>1</sup>	22.3 <sup>1</sup>	54.9 <sup>1</sup>
	Dry	N	N	N	N	N	N	N
Tuna fish	Fresh	N	0.90 <sup>6</sup>	N	0.76 <sup>6</sup>	75.24 <sup>6</sup>	22.37 <sup>6</sup>	N
	Dry	N	N	N	N	N	N	N
Blacktip shark	Fresh	N	1.15 <sup>6</sup>	N	1.56 <sup>6</sup>	77.78 <sup>6</sup>	20.04 <sup>6</sup>	N
	Dry	N	N	N	N	N	N	N

Data taken from <sup>1</sup>De Moreno *et al.* (2000); <sup>2</sup>Juárez-Luna (2012); <sup>3</sup>Sánchez *et al.* (2007); <sup>4</sup>Sadler (2003); <sup>5</sup>Manzi *et al.* 2004; <sup>6</sup>Castro-González *et al.* (2007); N= No data available.



### Comparison with other wild mushrooms

The comparison of proximal analysis results of *F. wolfeana* with wild mushrooms, are shown in Table 3, and data are shown in grams per 100 g of dry material. The average amount of protein in *F. wolfeana* is 29.83 g; *Amanita caesarea* had a higher value (34.77 g), while *Armillaria tabesceus* had a lower content (22.90 g). However, the contents of fat, carbohydrates and ash were the lowest for these two species. Therefore, we consider that *F. wolfeana* is a good diet alternative for people looking for foods low in fat and carbohydrates. This species is abundant in oak forests in Mexico, and large specimens grow there, which can weigh up to half a kilogram when fresh; that is why it is a highly prized species, because with a few basidiomata, they can cook for a family (six to eight members).

During the rainy season, *F. wolfeana* can replace other foods, such as meat, which in many cases is not accessible for most of the people of the communities. Also, this species could be dried and preserved for almost a year in a freezer.

Table 3. Basic comparison of *Fistulinella wolfeana* with other wild edible mushrooms (g/100g), data of dry material

Mushrooms species	Proteins	Fat	Carbohydrates	Ash
<i>Amanita caesarea</i>	<u>34.77</u>	3.50	55.63	6.05
<i>Armillaria mellea</i>	24.47	2.10	65.47	7.95
<i>Armillaria tabesceus</i>	22.90	2.54	<u>66.87</u>	7.63
<i>Boletus aureus</i>	27.17	4.47	62.10	6.25
<i>Cantharellus cibarius</i>	21.57	2.88	66.07	<u>9.44</u>
<i>Fistulina hepatica</i>	22.60	3.17	66.00	8.20
<i>Fistulinella wolfeana</i>	29.83	0.71	30.21	4.48
<i>Hygrophorus russula</i>	32.47	<u>6.00</u>	53.33	8.18
<i>Lepista nuda</i>	34.37	3.22	56.33	6.03
<i>Ramaria largentii</i>	28.80	5.67	58.87	6.67
<i>Russula delica</i>	26.10	2.88	66.07	<u>9.44</u>

Data taken from Ouzouni *et al.* (2009). The underlined number represents the highest value for each category.

### Antioxidant activity

Many common wild boletes are known to have a considerable antioxidant activity, comparable with our results (Figure 5), as indicated by several previous studies (Elmastas *et al.*, 2007; Ribeiro *et al.*, 2006; Sarikurkcu *et al.*, 2008; Tsai *et al.*, 2007). Looking for organic compounds profiles in boletes, we could find different values for fatty acids and phenolic compounds, that show high levels of linoleic, oleic and stearic acids (Kalaç, 2009). In general, mushrooms have low fat content, but in some cases, unsaturated acids such as oleic acid (C18:1) and linoleic acid (C18:2) are predominant over saturated fats (León-Guzmán *et al.*, 1997; Ribeiro *et al.*, 2009; Guillamón *et al.*, 2010). Some members of the Boletaceae family have shown the same predominance of unsaturated acids (Pedneault *et al.*, 2006). In this work, we report an increase in fat content when mushrooms were dried, but more studies are needed to determine the profile of fatty acids in raw, dry and even cooked mushrooms, in order to know if heat modifies the amount of fat. This could have implications as people dry them to consume them later.

It is important to notice that some species are considered health enhancers by people who consume them (Garibay-Orijel *et al.*, 2007; Alonso-Aguilar *et al.*, 2014), which makes sense, because it has been reported that the consumption of wild mushrooms play a role in the prevention of cardiovascular diseases (Guillamón *et al.*, 2010). Bautista (2013) published one of the most recent and complete reports about Mexican medicinal mushrooms traditionally used by Nahuas, Tseltales and Oto-




Figure 5. Percentage of radical scavenging activity with 2,2-diphenyl-1-picrylhydrazyl (DPPH) from methanol extracts diluted 1:10 and 1:100. Data were averaged from N=6, bars are standard deviation.

mies. The author cited a traditional medicine man, stating that: “not all the mushrooms are medicinal, but all edible mushrooms are potentially medicinal”. In the case of *F. wolfeana*, we could hypothesize that it has medicinal properties, but its use as one of the Mexican traditional wild mushrooms has not been reported. Elmastas *et al.* (2007) and Puttaraju *et al.* (2006) indicated that boletes contain phenolic compounds, suggesting that it is possible to find them in *Fistulinella* genus. In addition, many phenolic compounds have antioxidant potential (Ribeiro *et al.*, 2006). These components could also have the capacity to repel insects, as we have noticed in all the mushrooms collected for this work.

## CONCLUSIONS

We have reported here that *Fistulinella wolfeana* as a very common species in oak forests in Mexico, which produce abundant and robust basidiomata. We also described its important nutritional value and as a multifunctional non-timber resource that can be consumed when fresh, but also preserved and consumed after drying, a practice carried out in the three communities that comprised our study. Additionally, it is used as exchange for basic supplies, an activity that has been disappearing because of predominance of the modern monetary system in small villages. We reported a high antioxidant activity, but only using a 1:10 dilution, which decreases at higher dilutions and is considerably higher when not dissolved. Therefore, we can conclude that *F. wolfeana* is a good source of antioxidants. This work should encourage future studies on the bioactivity of Mexican members of the Boletaceae family..

## ACKNOWLEDGMENTS

We thank to CONACyT for the scholarship grant for postgraduate studies, Universidad Autónoma de Querétaro and PRODEP (SEP) for supporting this research. To all who helped in the ethnomycological work and especially to Sabina Wischin, who helped in determining the antioxidant activity and results interpretation.

## REFERENCES

- Alonso-Aguilar, L.E., A. Montoya, A. Kong, A. Estrada-Torres, R. Garibay-Orijel, 2014. The cultural significance of wild mushrooms in San Mateo Huexoyucan, Tlaxcala, Mexico. *Journal of Ethnobiology and Ethnomedicine* 10: 27.
- Bautista, J.A., 2013. Conocimiento tradicional de hongos medicinales en seis localidades diferentes del país. Tesis de licenciatura. Facultad de Ciencias, Universidad Nacional Autónoma de México, México.
- Bautista-Nava, E., Á. Moreno-Fuentes, 2009. Primer registro de *Calostoma cinnabarina* (Sclerodermatales) como especie comestible. *Revista Mexicana de Biodiversidad* 80: 561-564.
- Brand-Williams, W., M.E. Cuvelier, C. Berset, 1995. Use of a free radical method to evaluate antioxidant activity. *Wiss Technology* 28: 25-30.
- Burrola-Aguilar, C., O. Montiel, R. Garibay-Orijel, L. Zizumbo-Villareal, 2012. Conocimiento tradicional y aprovechamiento de los hongos comestibles silvestres en la región de Amanalco, Estado de México. *Revista Mexicana de Micología* 35: 1-16.
- Castro-González, M. I., V.A. Ojeda, B.S. Montaña, C.E. Ledesma, R.F. Pérez-Gil, 2007. Evaluación de los ácidos grasos n-3 de 18 especies de pescados marinos mexicanos como alimentos funcionales. *Archivos Latinoamericanos de Nutrición* 57: 85.
- De Moreno, L. A., A. Vidal, D. Huerta-Sánchez, Y. Navas, S. Uzcátegui-Bracho, N. Huerta-Leiden, 2000. Análisis comparativo proximal y de minerales entre carnes de iguana, pollo y res. *Archivos Latinoamericanos de Nutrición* 50: 409-415.
- De Román, M., E. Boa, S. Woodward, 2006. Wild-gathered fungi for health and rural livelihoods. *Proceedings of the Nutrition Society* 65: 190-197.
- Elmastas, M., O. Isildak, I. Turkecul, N. Temur, 2007. Determination of antioxidant activity and antioxidant compounds in wild edible mushrooms. *Journal of Food Composition and Analysis* 20: 337-345.
- Estrada-Torres A., R.M. Aroche, 1987. Acervo etnomicológico en tres localidades del municipio de Acambay, Estado de México. *Revista Mexicana de Micología* 3: 109-137.
- Ferreira, I. C., L. Barros, R. Abreu, 2009. Antioxidants in wild mushrooms. *Current Medicinal Chemistry* 16: 1543-1560.
- Food and Drug Administration (FDA), 2015. Sodium in Your Diet: Use the Nutrition Facts Label and Reduce Your Intake. Available In: <http://www.fda.gov/Food/IngredientsPackagingLabeling/LabelingNutrition/ucm315393.htm>
- García, J., 1999. Estudio sobre la taxonomía, ecología y distribución de algunos hongos de la familia Boletaceae (Basidiomycetos, Agaricales) en México. Tesis de Maestría inédita. Facultad de Ciencias Forestales, U.A.N.L.
- García, J., F. Garza-Ocañas, 2001. Conocimiento de los hongos de la familia Boletaceae de México. *Ciencia UANL* 4: 336-343.
- García J., D. Pedraza-Kamino, C.I. Silva-Barrón, R.L. Andrade-Melchor, J. Castillo-Tovar, 1998. Hongos del Estado de Querétaro. Universidad Autónoma de Querétaro, Querétaro.
- Garibay-Orijel, R., J. Cifuentes, 2004. Conocimiento Micológico Tradicional en la Planicie Costera del Golfo de México. *Revista Mexicana de Micología* 19: 57-70.
- Garibay-Orijel, R., J. Caballero, A. Estrada-Torres, J. Cifuentes. 2007. Understanding cultural significance, the edible mushrooms case. *Journal of Ethnobiology and Ethnomedicine* 3: 4.
- González, A., R. Valenzuela, 1993. Boletaceos y Gonfidaceos del Estado de México I. Discusiones sobre su distribución en diferentes tipos de vegetación, asociaciones ectomicorrizógenas, fenología y comestibilidad. *Revista Mexicana de Micología* 9: 35-46.
- Guillamón, E., A. García-Lafuente, M. Lozano, M.A. Rostagno, A. Villares, J.A. Martínez, 2010. Edible mushrooms: role in the prevention of cardiovascular diseases. *Fitoterapia* 81: 715-723.
- Instituto Nacional de Estadística Geografía e Informática (INEGI), 2009. Perfil sociodemográfico de la población que habla lengua indígena. México. Available In: [http://www.inegi.org.mx/prod\\_serv/contenidos/espanol/bvinegi/productos/censos/poblacion/poblacion\\_indigena/leng\\_indi/PHLI.pdf](http://www.inegi.org.mx/prod_serv/contenidos/espanol/bvinegi/productos/censos/poblacion/poblacion_indigena/leng_indi/PHLI.pdf)





- Juárez-Luna, M.A., 2012. Propiedades funcionales del hongo comestible *Pleurotus* y su relevancia en la alimentación regional. Tesis de maestría. Colegio de Postgraduados. Puebla. Mexico.
- Kalač, P., 2009. Chemical composition and nutritional value of European species of wild growing mushrooms: A review. *Food chemistry* 113: 9-16.
- Kirk, P.M., P.F. Cannon, J.C. David (eds.), 2008. *Ainsworth and Bisby's dictionary of the fungi*. 10th ed. CABI Bioscience, Oxon: 771.
- Konica Minolta Sensing, Inc. 2003. Comunicación precisa de los colores. Available on: <http://www2.konicaminolta.eu/Measuring/pcc/es/part1/index.html>
- Landeros, F., J. Castillo, G. Guzmán, J. Cifuentes, 2006. Los hongos (macromicetos) conocidos en el Cerro el Zamorano (Querétaro-Guanajuato), México. *Revista Mexicana de Micología* 22: 25-31.
- Lara-Vázquez, F., A.T. Romero-Contreras, C. Burrola-Aguilar, 2013. Conocimiento tradicional sobre los hongos silvestres en la comunidad Otomí de San Pedro Arriba; Temoaya, Estado de México. *Agricultura, Sociedad y Desarrollo* 10: 305-333.
- Manzi, P., A. Aguzzi, L. Pizzoferrato, 2001. Nutritional value of mushrooms widely consumed in Italy. *Food chemistry* 73: 321-325.
- Manzi, P., S. Marconi, A. Aguzzi, L. Pizzoferrato, 2004. Commercial mushrooms: nutritional quality and effect of cooking. *Food chemistry* 84: 201-206.
- Mapes, C., G. Guzmán, J. Caballero, 1981. *Etnomicología Purépecha: El conocimiento y uso de los hongos en la Cuenca del Lago de Pátzcuaro, Michoacán. Serie Etnociencias 2*. México. Dirección General de Culturas Populares, Secretaría de Educación Pública y Sociedad Mexicana de Micología e Instituto de Biología Universidad Nacional Autónoma de México: 79 p.
- Mata, G., 1987. Introducción a la etnomicología maya de Yucatán. El conocimiento de los hongos en Pixoy, Valladolid. *Revista Mexicana de Micología* 3: 175-188.
- Mohapatra, D., Z.M. Bira, J.P. Kerry, J.M. Frías, F.A. Rodrigues., 2010. Postharvest hardness and color evolution of white button mushrooms (*Agaricus bisporus*). *Journal of food science* 75: E146-E152.
- Mora, R.N., R. Valenzuela, 1997. Los macromicetos de la Sierra de Nanchititla, I. *Polibotánica* 5: 21-36.
- Morales, O., R. Medel, G. Guzmán, 2006. Primer registro de la comestibilidad de una especie de *Daldinia* (Ascomycota). *Revista Mexicana de Micología* 23: 103-105.
- Moreno-Fuentes, A., E. Aguirre-Acosta, L. Pérez-Ramírez, 2004. Conocimiento tradicional y científico de los hongos en el estado de Chihuahua, México. *Etnobiología* 4: 89-105.
- Ouzouni, P.K., P.G. Veltsistas, E.K. Paleologos, K.A. Riganakos, 2007. Determination of metal content in wild edible mushroom species from regions of Greece. *Journal of Food Composition and Analysis* 20: 480-486.
- Ouzouni, P.K., D. Petridis, W.D. Koller, K.A. Riganakos, 2009. Nutritional value and metal content of wild edible mushrooms collected from West Macedonia and Epirus, Greece. *Food Chemistry* 115: 1575-1580.
- Pedneault, K., P. Angers, A. Gosselin, R.J. Tweddell, 2006. Fatty acid composition of lipids from mushrooms belonging to the family Boletaceae. *Mycological Research* 110: 1179-1183.
- Pellicer-González, E., D. Martínez-Carrera, M. Sánchez, M. Aliphath, A. Estrada-Torres, 2002. Rural management and marketing of wild edible mushrooms in México. In: *International Conference on Mushroom Biology and Mushroom Products*. Cuernavaca, WSMBMP, Cuernavaca. Pp. 433-443.
- Pérez-Moreno, J., M. Martínez-Reyes, A. Yescas-Pérez, A. Delgado-Alvarado, B. Xoconostle-Cázares, 2008. Wild mushroom markets in central Mexico and a case study at Ozumba. *Economic Botany* 62: 425-436.
- Puttaraju, N.G., S.U. Venkateshaiah, S.M. Dharmesh, S.M.N. Urs, R. Somasundaram, 2006. Antioxidant activity of indigenous edible mushrooms. *Journal of Agricultural and Food Chemistry* 54: 9764-9772.
- Qi, L.L., M. Zhang, A.S. Mujumdar, X.Y. Meng, H.Z. Chen, 2014. Comparison of drying characteristics and quality of shiitake mushrooms (*Lentinus edodes*) using different drying methods. *Drying Technology* 32: 1751-1761.
- Ribeiro, B., J. Rangel, P. Valentão, P. Baptista, R.M. Seabra, P.B. Andrade, 2006. Contents of carboxylic acids and two phenolics and antioxidant activity of dried Portuguese wild edible mushrooms. *Journal of Agricultural and Food Chemistry* 54: 8530-8537.
- Ribeiro, B., P.G. de Pinho, P.B. Andrade, P. Baptista, P. Valentão, 2009. Fatty acid composition of wild edible mushrooms species: A comparative study. *Microchemical Journal* 93: 29-35.
- Rodríguez, O., M. Herrera-Fonseca, M.D.R. Sánchez-Jácome, I. Álvarez, R. Valenzuela, J. García, L. Guzmán-Dávalos, 2010. Catálogo de la micobiota del bosque La Primavera, Jalisco. *Revista Mexicana de Micología* 32: 29-40.
- Sadler, M., 2003. Nutritional properties of edible fungi. *Nutrition Bulletin* 28: 305-308.
- Sarikurkcü, C., B. Tepe, M. Yamac, 2008. Evaluation of the antioxidant activity of four edible mushrooms from the Central Anatolia, Eskisehir-Turkey: *Lactarius deterrimus*, *Suillus collinitus*, *Boletus edulis*, *Xerocomus chrysenteron*. *Bioresource Technology* 99: 6651-6655.
- Singer, R., J. García, L.D. Gómez-Pignataro, 1991. The Boletineae of México and Central America III. Los Boletineae de México y Centroamérica III. *Nova Hedwigia* 102: 1-97.
- Ruan-Soto, F., 2007. 50 años de etnomicología en México. *Lacandonia* 1: 97-108.
- Tsai, S.Y., H.L. Tsai, J.L. Mau, 2007. Antioxidant properties of *Agaricus blazei*, *Agrocybe cylindracea*, and *Boletus edulis*. *LWT-Food Science and Technology* 40: 1392-1402.
- U.S. Food and Drugs Agency (FDA). 2015. Sodium in Your Diet: Using the Nutrition Facts Label to Reduce Your Intake. Available In: <http://www.fda.gov/Food/IngredientsPackagingLabeling/LabelingNutrition/ucm315393.htm>.
- Villanueva, K.T., G.B. Williams, M. Herrera-Fonseca, A. Galván-Corona, O.R. Alcántar, 2008. Avances en el conocimiento de los hongos macromicetos del Cerro El Venado, Mezcala, Mpio. de Poncitlán, Jalisco. Avances en la Investigación Científica en el CUCBA. XIX Semana Nacional de Investigación Científica, Universidad de Guadalajara, Guadalajara. Pp. 223-230.
- Wu, G., B. Feng, J. Xu, X.T. Zhu, Y.C. Li, N.K. Zeng, Z.L. Yang, 2014. Molecular phylogenetic analyses redefine seven major clades and reveal 22 new generic clades in the fungal family Boletaceae. *Fungal Diversity* 69: 93-115.
- Zamudio, S., J. Rzedowski, W.E. Carranza, G. Claderon, 1992. La Vegetación del estado de Querétaro: panorama preliminar. Instituto de Ecología, Pátzcuaro.